

## Coping with Intrusive Thoughts after a Traumatic Loss

### What are intrusive thoughts?

- Intrusive thoughts are upsetting memories of a traumatic event that enter a survivor's mind even when they are trying not to think about it.
- Survivors may feel the same upsetting and overwhelming reactions they had when the event occurred.
- Survivors often report that intrusive thoughts are some of the most difficult trauma reactions to manage, but there are steps you can take to decrease their frequency and intensity.

### Here are some techniques you can use to gain control over intrusive thoughts:

#### 1. Notice

- If you notice that you are having an intrusive thought, consciously ask yourself, "Is there anything I will gain from recalling this moment?"
- Give yourself permission not to relive that moment.
- Remind yourself that your loved one deserves to be honored for who they were and what they contributed when they lived. You don't need to focus on the circumstances of their death.

#### 2. Breathe

- Controlled breathing has a physiological effect on your body that slows the heart rate and allows the brain to refocus on the sensation of taking slow breaths.
- If you experience anxiety, sadness, anger, or any other distressing emotions associated with an intrusive thought, you can try to ease it by slowing down your breathing. Try:
  - a. Getting in a comfortable position;
  - b. Inhaling slowly through your nose for four seconds;
  - c. Pausing and holding your breath for about four seconds;
  - d. Exhaling slowly through your mouth for about five seconds;
  - e. Breathing from your stomach instead of your chest (deep breaths should move your abdomen area in and out); and
  - f. Listening to your breathing and keeping it as even as possible.

#### 3. Replace

- Again, you don't need to focus on the distressing events surrounding the loss of your loved one.
- Here are some ways you can replace your distressing thoughts with comforting ones:
  - a. Allow yourself to recall a special or happy memory with your loved one.
  - b. If you are comforted by relaxing imagery, imagine a place that feels peaceful to you.
  - c. Try refocusing on literally any other topic; the topic is less important than the process of giving the brain another thought to focus on. Here are some examples:

1. Think about the top five sports moments in history;
2. Imagine the design of a beautiful dress; or
3. Recall your favorite episode from your favorite TV show.

#### 4. Redirect

- Your senses can be very helpful in refocusing your mind on the present moment.
- Use “grounding techniques” to reorient yourself with the current moment:
  - a. Splash cold water on your face or hold an ice cube in your hand.
  - b. Drink something hot or cold.
  - c. Notice colors or objects around the room. Count how many blue items you see.
  - d. Smell something with an aroma you enjoy, such as a scented candle, a food, or a bathroom product.
  - e. Listen to the sounds of the environment around you. Can you hear the air conditioner or heater? Do you hear traffic in the distance?
  - f. Pay attention to how your body feels. Are you warm? Cold? Are your shoes comfortable? Are you comfortable in the position you are sitting in?

#### 5. Reflect

- Intrusive thoughts are sometimes triggered by something happening in your environment. Becoming aware of potential triggers can give you more control over your thoughts, even if you can't avoid the triggers entirely.
- When you feel calm, reflect on what might be causing your intrusive thoughts:
  - a. Are there certain times of the day that these thoughts seem most likely to occur?
  - b. Are the memories of the event linked to the people you are with, or linked to the place where you currently are?
  - c. Is there a sensory component to the thought you are experiencing? Are certain smells, sights, or sounds possible triggers?
  - d. Is there anything going on in your environment that could remind you of aspects of the event? Are people discussing a certain topic? Are there certain materials on TV or social media?

#### References

- Berkowitz, S., & Marans, S. (2011). *The Child and Family Traumatic Stress Intervention: Implementation Guide for Providers*.
- Lynch, J., Mack, L., Benesek, J., Marshall, C., Cevinger, L., McHenry, S., Reynolds, S., Mutchler, B., Meyer, B., Panissidi, D., Jones, A., & Hall, S. (2015). *PTSD recovery program treatment manual*. Retrieved from [https://www.mirecc.va.gov/docs/vsn6/PTSD\\_Recovery\\_Group-Client\\_Manual\\_3rd\\_edition.pdf](https://www.mirecc.va.gov/docs/vsn6/PTSD_Recovery_Group-Client_Manual_3rd_edition.pdf)
- Sanderson, C. (2009). Introduction to counselling survivors of interpersonal trauma. London: Jessica Kingsley Publishers.

The content of this product was produced by FrontLine Service under Grant Number 2016-XV-GX-K019, awarded by the Office for Victims of Crime, Office of Justice Programs, U.S. Department of Justice. The design was created by The University of Akron with funding from the Peg's Foundation. The opinions, findings, and conclusions or recommendations expressed in this product are those of the contributors and do not necessarily represent the official position or policies of the U.S. Department of Justice, The University of Akron or the Peg's Foundation.